

Do common buzzards *Buteo buteo* migrate across the Channel of Sicily during autumn?

NICOLANTONIO AGOSTINI¹, PAOLO AMATO², ANTONINO PROVENZA², MICHELE PANUCCIO³

¹ via Carlo Alberto 4, I-89046 Marina di Gioiosa Jonica (RC), (nicolantonio@tiscalinet.it); ² LIPU sez. di Alcamo, via G. D'Arco 2, I-91011 Alcamo (TP); ³ MEDRAPTORS (Mediterranean Raptor Migration Network), via M. Fioretti 18, I-00152 Roma

While Mediterranean populations of the common buzzard *Buteo buteo* are sedentary, those breeding in central, eastern and northern Europe are summer resident or partially migratory. During autumn migration, tens of thousands of birds are recorded both at Falsterbo (Sweden) and at the Bosphorus, while lower numbers cross the Strait of Gibraltar (Cramp and Simmons 1980). In the central Mediterranean region, observations made in southern Italy and Malta both during autumn and spring migration showed that very few buzzards move in this area (Agostini 2001, Agostini and Logozzo 1995, 1997, 1998, Agostini and Malara 1997, Agostini and Panuccio 2003, Agostini *et al.* 2000, 2004, Corso 2001, Guglielmi *et al.* 2003, Jonzén and Pettersson 1999, Panuccio *et al.* 2004, Sammut and Bonavia 2004). Since these raptors mostly use soaring flight over land during migration, thus avoiding long sea crossings (Kerlinger 1989), these results are not unexpected. However, observations on the post-reproductive migration of this species across the Channel of Sicily during the peak passage (late September - mid October) are lacking (Cramp and Simmons 1980). In this note, we provide data on buzzard migration recorded at the island of Marettimo between 15-30 September 2000 and 3-18 October 2002.

Study area and methods - This island is located about 30 km off western Sicily and 130 km NE of the Cap Bon Peninsula (Tunisia). Observations, aided with binoculars and telescopes, were made using an observation post at the altitude of c 500 m. At this site thousands of raptors, mostly black kites *Milvus migrans* and adult honey buzzards *Pernis apivorus*, concentrate during autumn migration between the end of August and mid September (Agostini *et al.* 2000, 2004).

Results and discussion - During the second half of September 2000, a total of 332 raptors was counted, mostly marsh harriers *Circus aeruginosus*, while only 4 (1.2 %) buzzards were reported. Not a single buzzard was observed between 3 and 18 October 2002. In this period, the short-toed eagle *Circaetus gallicus* was the most frequent species (Tab. 1). Finally, 6 buzzards were reported over the island during systematic observations made between 1-31 October 2004 (Amato *et al.* pers. obs.). Although we cannot exclude that some buzzards crossed the Channel of Sicily bypassing Marettimo (Agostini *et al.* 2004), these data agree with the conclusion that a true migratory flow across the Central Mediterranean is non-existent (Agostini 2002). However,

late and/or irregular movements of birds belonging to the Sicilian population (perhaps depending on ecological conditions in their breeding grounds) and/or coming from eastern Europe, could explain the existence of the small population of this species wintering in northern Tunisia such as the spring passage of some steppe buzzards *Buteo buteo vulpinus* at the Strait of Messina. In this framework, counts concerning hundreds of buzzards recorded at the Cap Bon promontory during spring 1974 and 1975 (Thiollay 1977) were probably due to repeated counting of birds belonging to the population wintering in Tunisia.

References - Agostini N 2001. *Buteo* 12: 99-102 • Agostini N 2002. In: *Manuale di Ornitologia*, Edagricole-II Sole 24 Ore, Bologna • Agostini N, Logozzo D 1995. *Riv. ital. Orn.* 64: 117-120 • Agostini N, Logozzo D 1997. *Avocetta* 21: 174-179 • Agostini N, Logozzo D 1998. *Riv. ital. Orn.* 68: 153-157 • Agostini N *et al.* 2000. *Avocetta* 24: 95-99 • Agostini N, Malara G 1997. *Riv. ital. Orn.* 66: 174-176 • Agostini N, Panuccio M 2003. *Riv. ital. Orn.* 73: 165-167 • Agostini N *et al.* 2004. *Ring* 26: 71-78 • Corso A 2001. *British Birds* 94: 196-202 • Cramp S, Simmons KEL 1980. Oxford Univ. Press, Oxford • Guglielmi R *et al.* 2003. *Avocetta* 27: 69 • Jonzén N, Pettersson J 1999. *Avocetta* 23: 65-72 • Kerlinger P 1989. Univ. Chicago Press, Chicago • Panuccio M *et al.* 2004. *British Birds* 97: 400-403 • Sammut M, Bonavia E 2004. *British Birds* 97: 318-322 • Thiollay JM 1977. *Alauda* 45: 115-121.

Table 1 – Raptors observed over the island of Marettimo between 15-30 September 2000 and 3-18 October 2002.

Species	15-30 September	3-18 October
<i>Circaetus gallicus</i>	18	125
<i>Circus aeruginosus</i>	220	27
<i>Pernis apivorus</i>	45	3
<i>Milvus migrans</i>	15	4
<i>Buteo buteo</i>	4	-
<i>Hieraaetus pennatus</i>	9	2
<i>Neophron percnopterus</i>	2	-
<i>Aquila sp.</i>	2	-
<i>Accipiter nisus</i>	4	-
<i>Falco subbuteo</i>	5	-
<i>Falco tinnunculus</i>	4	-
<i>Falco naumanni</i>	1	-
<i>F. tinnunculus/naumanni</i>	3	1
<i>Falco biarmicus</i>	-	1